

proportional hazards regression analysis identified FA score was an independent predictor for OS and CSS.

**Conclusions:** The evaluation of preoperative FA score can be regarded as an independent prognostic factor for predicting OS and CSS of patients with UTUC. The fibrinogen and albumin levels are low cost, routine measured and easy accessibility in clinical practice.

**Keywords:** Urinary tract urothelial carcinoma (UTUC); plasma fibrinogen; serum albumin; prognosis

doi: 10.21037/tau.2017.s049

**Cite this abstract as:** Cui J, Zhang Z, Jiang X, Du J, Liu Y, Zhao H, Shi B. Prognostic scores based on the preoperative plasma fibrinogen and serum albumin level as a prognostic factor in patients with upper urinary tract urothelial carcinoma. *Transl Androl Urol* 2017;6(Suppl 3):AB049. doi: 10.21037/tau.2017.s049

## AB050. Prognostic relevance of preoperative platelet-to-albumin ratio in patients with upper urinary tract urothelial carcinoma

Hongda Zhao, Jianfeng Cui, Yan Li, Meng Yu, Nin Zhang, Shiyu Wang, Benkang Shi

Department of Urology, Qilu Hospital of Shandong University, Jinan 250012, China

**Background:** The aim of this study is to assess the value of platelet-to-albumin ratio on overall survival (OS) and cancer specific survival (CSS) in patients with upper urinary tract urothelial carcinoma (UTUC).

**Methods:** We retrospectively evaluated 169 patients' clinicopathological data. The Kaplan-Meier analysis was used to assess the OS and CSS and test the equivalences of the curves by log-rank tests. Univariate and multivariate survival analyses were carried out using Cox proportional hazards regression model.

**Results:** Platelet-to-albumin ratio (PAR) was significantly associated with OS and CSS. Multivariate analyses revealed that PAR was an independent prognostic factor in patients with UTUC. Patients with high PAR value had a higher probability of OS and CSS, compared to patients with low PAR.

**Conclusions:** The platelet and serum albumin level are low cost and easy to obtain, PAR may be a useful independent prognostic factor in patients with UTUC.

**Keywords:** Urinary tract urothelial carcinoma (UTUC); platelet; albumin; prognosis

doi: 10.21037/tau.2017.s050

**Cite this abstract as:** Zhao H, Cui J, Li Y, Yu M, Zhang N, Wang S, Shi B. Prognostic relevance of preoperative platelet-to-albumin ratio in patients with upper urinary tract urothelial carcinoma. *Transl Androl Urol* 2017;6(Suppl 3):AB050. doi: 10.21037/tau.2017.s050

## AB051. Simultaneous bilateral retroperitoneal laparoscopic nephron sparing surgery: a case report and evaluation of the technique

Ming Tong

Department of Urology, The First Affiliated Hospital of Jinzhou Medical University, Jinzhou 121000, China

**Background:** Bilateral renal cell carcinoma (RCC) is an extremely rare renal tumor, which even more infrequently presents with two different types of pathology.

**Methods:** Bilateral RCC is an extremely rare renal tumor, accounting for 3–4% of sporadic RCC. Bilateral and multifocal RCC are usually of a single histological type and infrequently of different types.

**Results:** Currently, nephron-sparing surgery (NSS) has become the standard for treating bilateral RCC, even for tumors greater than 4 cm. Laparoscopic NSS has developed

as a minimally invasive procedure and novel technique for the treatment of RCC.

**Conclusions:** To our knowledge, this is the first case to date of bilateral RCC treated with simultaneous retroperitoneal laparoscopic nephron-sparing surgery (RLNSS). Here we indicate the feasibility of this management and discuss the advantages and disadvantages of this technique.

**Keywords:** Renal cell carcinoma (RCC); nephron-sparing surgery (NSS); simultaneous bilateral

doi: 10.21037/tau.2017.s051

**Cite this abstract as:** Tong M. Simultaneous bilateral retroperitoneal laparoscopic nephron sparing surgery: a case report and evaluation of the technique. *Transl Androl Urol* 2017;6(Suppl 3):AB051. doi: 10.21037/tau.2017.s051

## AB052. ZEB1 promotes vasculogenic mimicry formation in prostate cancer is associated with epithelial-mesenchymal transition

Hua Wang, Zongren Wang, Shaopeng Qiu

Department of Urology, The First Affiliated Hospital, Sun Yat-sen University, Guangzhou 510080, China

**Background:** This study investigated the role of ZEB1 in vasculogenic mimicry (VM) formation and the interplay between VM and epithelial-mesenchymal transition (EMT).

**Methods:** Ninety-two prostate cancer tissue specimens were stained by CD34 and periodic acid Schiff. Then, we stained ZEB1 protein in the consecutive sections. Moreover, prostate cancer cells were subjected to ZEB1 knockdown using ZEB1 siRNA and then to 3D culture assay. EMT related maker was also evaluated.

**Results:** The data showed that the presence of VM and high ZEB1 expression were associated with higher Gleason score, TNM stage, and lymph node and distant metastases. ZEB1 knockdown reduced VM formation and the

expression of EMT-related in prostate cancer cells.

**Conclusions:** In the current study was the first to reveal that ZEB1 played an important role in VM formation in prostate cancer *ex vivo* and *in vitro*. Mechanistically, this process may have a relationship with EMT.

**Keywords:** Prostate cancer; vasculogenic mimicry (VM); ZEB1, epithelial-mesenchymal transition (EMT)

doi: 10.21037/tau.2017.s052

**Cite this abstract as:** Wang H, Wang Z, Qiu S. ZEB1 promotes vasculogenic mimicry formation in prostate cancer is associated with epithelial-mesenchymal transition. *Transl Androl Urol* 2017;6(Suppl 3):AB052. doi: 10.21037/tau.2017.s052

## AB053. Labeling of prostate tumor-specific replication-selective oncolytic adenoviruses with radioactive 125I: inhibitory effects on prostate cancer cell

Lin Hao, Conghui Han

Department of Urology, Xuzhou Central Hospital, Xuzhou 221000, China

**Background:** The authors established a 125I-labeled replication-selective oncolytic adenovirus and human telomerase reverse transcriptase/prostate-specific antigen (125I-RSOAds-hTERT/PSA) oncolytic adenovirus marker and investigated the effects of different labeling conditions. This study also explored the possible mechanism whereby 125I-RSOAds-hTERT/PSA inhibited the proliferation of prostate cancer cells.

**Methods:** N-bromosuccinimide (NBS) was used as an oxidant for 125I labeling, and various concentrations of oncolytic viruses and NBS were prepared to determine the optimal conditions for labeling. The effects of the