



# UroVysion fluorescence in situ hybridization in urothelial carcinoma: a narrative review and future perspectives

Takashi Nagai<sup>1</sup>, Taku Naiki<sup>1^</sup>, Toshiki Etani<sup>1</sup>, Keitaro Iida<sup>1</sup>, Yusuke Noda<sup>1</sup>, Nobuhiko Shimizu<sup>1</sup>, Teruki Isobe<sup>1</sup>, Satoshi Nozaki<sup>1</sup>, Takehiko Okamura<sup>2</sup>, Ryosuke Ando<sup>1</sup>, Noriyasu Kawai<sup>1</sup>, Takahiro Yasui<sup>1</sup>

<sup>1</sup>Department of Nephro-urology, Nagoya City University, Graduate School of Medical Sciences, Nagoya, Japan; <sup>2</sup>Department of Urology, Anjo Kosei Hospital, Anjo City, Japan

*Contributions:* (I) Conception and design: T Nagai, T Naiki, T Okamura, T Yasui; (II) Administrative support: All authors; (III) Provision of study materials or patients: T Etani, K Iida; (IV) Collection and assembly of data: Y Noda, N Shimizu, T Isobe, S Nozaki; (V) Data analysis and interpretation: R Ando, N Kawai; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

*Correspondence to:* Taku Naiki, MD, PhD. Department of Nephro-urology, Nagoya City University, Graduate School of Medical Sciences, Kawasumi 1, Mizuho-cho, Mizuho-ku 467-8601, Nagoya, Japan. Email: naiki@med.nagoya-cu.ac.jp.

**Abstract:** The number of patients with urothelial carcinoma (UC) is high, with a corresponding demand for detecting UC easily and non-invasively. Cystoscopy and urine cytology, with widely known diagnostic accuracies, are the gold standards for identifying UC originating from the bladder. However, cystoscopy or other tests, such as ureteroscopy or retrograde pyelography, are uncomfortable for patients. Tests for urinary biomarkers are expected to satisfy the demand for less invasive tests that will benefit patients with anxiety for invasive tests such as cystoscopy or ureteroscopy. Although several urinary biomarkers have been reported to support the diagnosis or follow-up of UC, their use in the clinic is uncommon. The UroVysion test examines urinary biomarkers using a multitarget, multicolor fluorescence in situ hybridization (FISH) assay. The test uses exfoliated cells found in urine and is a mixture of centromeric fluorescent denatured chromosome enumeration probes for chromosomes 3, 7, and 17 (labelled stratum red, spectrum green and spectrum aqua, respectively), and a locus-specific identifier probe for 9p21 (spectrum gold). It is used for the initial diagnosis of patients with hematuria or the monitoring of patients previously diagnosed with bladder cancer. Almost 20 years have passed since UroVysion was approved by the U.S. Food and Drug Administration, and so this is a well-established test. However, room exists for further research, with numerous reports on this test having been recently published. In order to update our knowledge, we herein present a brief overview of UroVysion and its features that follows the latest findings as they relate to UC.

**Keywords:** UroVysion; urothelial cancer; fluorescence in situ hybridization (FISH)

Submitted Aug 28, 2020. Accepted for publication Dec 08, 2020.

doi: 10.21037/tau-20-1207

**View this article at:** <http://dx.doi.org/10.21037/tau-20-1207>

## Introduction

Urothelial carcinoma (UC) develops in either the upper (pyelocaliceal cavities and ureter) or lower (bladder and urethra) portions of the urinary tract. As the 10th most common cancer worldwide, approximately 550,000 cases of bladder cancer (BCa) are newly diagnosed annually (1).

Although upper urinary tract carcinoma is an uncommon carcinoma, it accounts for 5–10% of all UCs (2). Generally, patients with UC complain of gross hematuria. The standard diagnostic tools available for the detection of BCa are cystoscopy, ureteroscopy, ultrasonography, and urine cytology. Magnetic resonance imaging and

<sup>^</sup> ORCID: 0000-0002-7638-6048.

Fluorophore	Chromosome	Cytogenic location	Probe name
	3	3p11.1-q11.1 Alpha Satellite DNA	Vysis CEP3 SpectrumRed
	7	7p11.1-q11.1 Alpha Satellite DNA	Vysis CEP7 SpectrumGreen
	9	9p21	Vysis LSI p16 (9p21) SpectrumGold Probe
	17	17p11.1-q11.1 Alpha Satellite DNA	CEP 17 SpectrumAqua

**Figure 1** Details of fluorescence in situ hybridization (FISH) probes.

computed tomography are also used for staging. Retrograde pyelography is used for detecting ureteral or renal pelvic UCs. However, cystoscopy, including retrograde pyelography, is reported as being uncomfortable for patients (3). Moreover, it is difficult to detect small, flat carcinomas using the aforementioned techniques. For the differential diagnosis of hematuria and the follow-up of urothelial cancers, a non-invasive, highly accurate test is therefore required.

The measurement of urinary biomarkers is one method of examining patients for UCs. Its simplicity of use and non-invasiveness meet the needs of both physicians and patients. To date, tests involving several urinary biomarkers are in clinical use: CxBladder (Pacific edge Ltd., Dunedin, New Zealand), bladder tumor-associated antigen (BTA; Polymedco Inc., Cortlandt Manor, NY, USA), Immnocytt/uCyt+ (Diagnocure, Inc., Quebec City, Canada), nuclear matrix protein 22 (NMP22; Matritech. Inc., Newton, MA, USA), AssureMDx (MDxHealth, Irvine, CA, USA), and UroVysion (Abbott Molecular Inc., Des Plaines, IL, USA). These have all been approved by the U.S. Food and Drug Administration (FDA) as non-invasive tests to diagnose BCa (4). They have also supported the diagnosis of UC or are used for follow-up (5).




The UroVysion test for urinary biomarkers is made up of a multitarget, multicolor fluorescence *in situ* hybridization (FISH) assay (6). One of the features of this test is its dependence on specific chromosomal abnormalities occurring in high numbers in urothelial cancers (7). The use of several probes in this assay means an improvement in sensitivity compared to using single probes (6). The test is performed on exfoliated cells in urine using centromeric fluorescent denatured chromosome enumeration probes for chromosomes 3, 7, and 17 (stratum red, spectrum green, and spectrum aqua, respectively), and a locus-specific identifier probe for 9p21 (spectrum gold), with a high sensitivity for detecting carcinomas (8). The details of each probe are outlined in *Figure 1*. Indications for UroVysion analysis

in urinary cytology include: atypical urinary cytology, for control after intravesical Bacillus Calmette-Guérin (BCG) treatment, atypical upper urinary tract cytology, follow-up after transurethral resection, and hematuria in patients with increased risk of UC (9). This test was approved by the FDA in 2001 and has been used to diagnose the recurrence of BCa from 2001 and to examine gross hematuria from 2005. In addition, it has been suggested that UroVysion FISH be used to judge the response to intravesical BCG therapy and in adjudicated equivocal cytology in American Urological Association guidelines (level of recommendation: expert opinion). UroVysion is also described as a potential urinary biomarker for the detection of invisible tumors in European Association of Urology guidelines. In National Comprehensive Cancer Network guidelines, UroVysion or NMP22 are considered markers useful in monitoring the recurrence of BCa. However, though the various guidelines described above highlight the usefulness of UroVysion, it is necessary to recognize variations in its use with regard to clinical settings.

Indeed, the use of UroVysion have not changed significantly in recent years. However, the treatment of UC has gradually improved due to the appearance of new therapies such as molecular-targeted drugs or a combination of intravesical therapy and immune-oncology drugs. Genetic mutations are therapeutic targets and screening for these seems to be required to achieve tailor-made therapy. A method that detects genetic mutations in urine samples using FISH can change the future treatment of UC. We present the following article in accordance with the Narrative Review reporting checklist (available at <http://dx.doi.org/10.21037/tau-20-1207>).

## Methods

A literature review in English was conducted using PubMed

		Negative	Sample of positive pattern	
				
FISH probe		Number of signals		
CEP3		2	3	2
CEP7		2	3	3
LSI 9p21		2	0	0
CEP17		2	3	2
Result of UroVysion				
Aneuploidy			✓	
Deletion			✓	✓

**Figure 2** Patterns of fluorescence in situ hybridization (FISH) signals.

and Google Scholar databases up to September 2020. Search terms included: “UroVysion” AND “urology” or “FISH” AND “urology”. The studies were reviewed by three authors to assess whether they were appropriate.

### What is FISH and UroVysion?

Genetic abnormalities, including mutations and chromosome and gene copy number irregularities, drive the development and progression of cancer (9). FISH is a method to detect genetic alterations by hybridizing such genes with oligonucleotide probes labeled with a fluorescent substance or an enzyme to identify the target using a fluorescence microscope. FISH is widely used for detecting chromosomal abnormalities in various cancers. Moreover, it is thought to be an objective examination since it clearly detects a specific gene locus compared to conventional pathological examinations.

UroVysion is a molecular cytology test for BCa that uses FISH for the detection of aneuploidy in chromosomes 3, 7, and 17, and the loss of the 9p21 locus. The four probes are used in response to the increased copy numbers of chromosomes 3, 7, and 17 found in BCa as well as the deletion of 9p21, where *P16*, the tumor suppressor gene, resides (9). UroVysion is assessed by measuring the number of fluorescent signals in at least 25 cells that show an

abnormal morphology. According to the manufacturer’s criteria, the chromosome pattern needs to be recorded only if (I) there is a gain (i.e., three or more signals) of more than two of chromosomes 3, 7, or 17, or (II) if both copies of LSI 9p21 are absent. The test is positive if there has been an increase of four or more cells showing multiple chromosomes or more than 12 cells that have lost a 9p21 signal. The negative signal pattern and some of positive signal patterns are shown in *Figure 2*.

Although it appears to be an objective examination compared to urine cytology, UroVysion results sometimes depend on the skill of the laboratory technicians who perform the test. Moreover, UroVysion analysis should be conducted by well-experienced laboratory technicians in cytopathology (10). This is because the interpretation of UroVysion may be difficult due to the use of fluorescence (6). Physicians should also interpret UroVysion results with regard to clinical findings and not leave examinations to laboratory technicians.

Surveillance with the UroVysion can provide prognostic information for patients with non-muscle invasive UC (11); however, it is crucial to consider the patient’s background in order to attain an accurate diagnosis. Results may vary with regard to sensitivity and specificity, which might be due to selection bias of patients. UroVysion and urine cytology are not thought to be affected by the presence of erythrocytes or leukocytes compared to the NMP22 test (12). However,

the UroVysion test yields false positives due to the presence of umbrella cells, as suggested by the presence of chromosome tetraploidy, or heteroploidy, which appears because of human polyomavirus infection (13-15).

Most papers have compared the UroVysion test to urine cytology or other urinary biomarkers. However, comparisons are difficult simply because of the variations in clinical settings. We have reviewed previous reports based on each clinical setting.

### Screening hematuria

Hematuria is one of the major complaints of UC including BCa. Generally, urine cytology is conducted to diagnose urinary tract cancers or in a work-up of hematuria. The definitive role of urinary biomarkers is yet to be established. However, a recent meta-analysis of urinary biomarkers for the evaluation of primary hematuria concluded that evidence for the use of urinary biomarkers was limited with regard to diagnostic performance, casting doubt on the routine use of these instead of cystoscopy (4).

Few reports exist regarding the evaluation of hematuria by UroVysion. UroVysion has been described as an accurate, more sensitive test than voided cytology, and has been used to detect BCa in patients showing gross or microscopic hematuria of all stages and grades (16). The UroVysion test detected 69% of UCs while cytology detected 38%; the gap between these widened if low-grade, low-stage tumors were excluded (16). Based on these data, the FDA approved UroVysion for examining hematuria from 2005.

### BCa

Lists of major studies that compared UroVysion and urine cytology are shown in *Table 1*.

In a meta-analysis, Hajdinjak (2008) reported the superiority of UroVysion to urine cytology, showing a sensitivity of 72% and specificity of 84% compared with 42% and 96% for cytology, respectively (24). However, such differences disappeared when superficial cases were excluded (24). Several conflicting reports exist with regard to the diagnosing power of UroVysion. In a surveillance of non-muscle invasive BCa, UroVysion did not yield urine cytology for the detection of visible BCa (25). In Japanese reports, a similar study reported that UroVysion alone was insufficient for detecting BCa (26).

Combining UroVysion and urine cytology was reported

as a method to improve sensitivity (25,26). Daniely *et al.* (2007) reported on the utility of urine cytology and UroVysion by using an automated microscope (27). Kojima *et al.* (2018) described the usefulness of a consecutive UroVysion test: Two consecutive positive UroVysion test results led to a cancer detection rate of 14.8%, which was greater than a positive result in either (7.2%) or neither (1.2%) of the two tests for patients at a 3-month follow-up (28).

However, several clinical situations have led to suspicious urine cytology results. Ferra *et al.* (2009) reported that a negative UroVysion result did not exclude the presence of high-grade UC (29). However, a recent prospective study showed the utility of UroVysion in a comparison to urine cytology in terms of sensitivity and specificity, and recommended the use of UroVysion in case of a suspicious result in urine cytology (30).

The interpretation of a 9p21 loss differs in several reports. Zellweger (2006) reported that the presence of a 9p21 deletion was significantly associated with recurrence (13). However, in a prospective study of 1,595 male chemical workers, a loss of 9p21 was less predictive for detecting BCa because of a clinical setting in which few low-grade cancers existed (31).

### UC of upper urinary tract

Ureteroscopy is a major tool for the diagnosis of upper tract UC. However, concerns exist because of its invasiveness, which may lead to severe complications such as bleeding, ureteral perforation, or stripping of the ureter (32). Compared to those of urine cytology, the sensitivity and specificity of UroVysion was significantly higher (76.7% *vs.* 36% and 94.7% *vs.* 100%, respectively (33). A multicentre cohort study found a higher diagnosing accuracy for UroVysion in UC (34). The overall sensitivity of UroVysion was higher than that of the cytology (100% *vs.* 20.8%), with the specificity 89.5% for UroVysion and 97.4% for cytology (34). Freund *et al.* (2019) reported that using UroVysion for 1 mL of passively collected upper tract urine in detecting upper urinary tract UC was feasible, even if the sample size used was small (35). Compared to negative results, positive FISH results predicted more advanced tumor stages and higher tumor grades (36). In addition, UroVysion was also reported to improve diagnostic accuracy for UC of the upper urinary tract when combined with urine cytology (37,38).

Conflicting opinions exist with regard to using UroVysion for the surveillance of upper tract UC. Fernández *et al.* (2012) examined the utility of UroVysion

Table 1 Major studies that compared UroVysion and cytology

Authors	Year	Number of samples	Number of Patients	Eligible patients	Number of patients with cancer	Probe	Positive definition	Overall sensitivity	Overall specificity
Halling <i>et al.</i> (17)	2000	280	265	165 patients with UC history and 115 patients with genitourinary symptoms	75	3, 7, 17, 9p21	5 or more cells with gains of 2 or more chromosomes	UroVysion: 81% vs. cytology: 58%	UroVysion: 96% vs. cytology: 98%
Sarosdy <i>et al.</i> (18)	2002	176, 275	176, 275	176 patients with UC in the previous 9 months and 275 volunteers without previous UC	62	3, 7, 17, 9p21	Product labeling	UroVysion: 71% vs. BTA stat, cytology 50%, 26%	UroVysion: 94.5
Laudadio <i>et al.</i> (19)	2005	521	300	Over a 1-year period consecutive cases being evaluated for bladder cancer recurrence or a potential new diagnosis of UC	44	3, 7, 17, 9p21	Product labeling	UroVysion: 73% vs. cytology: 34%	UroVysion: 65% vs. cytology: 93%
Sarosdy <i>et al.</i> (16)	2006	473	497	Patients with gross or microscopic hematuria	51	3, 7, 17, 9p21	Product labeling	UroVysion: 69% vs. cytology: 38%	UroVysion: 78%
Caraway <i>et al.</i> (20)	2010	941	600	Patients who were monitored for recurrent UC or evaluated for urinary symptoms	170	3, 7, 17, 9p21	Product labeling	UroVysion: 58% vs. cytology: 39%	UroVysion: 66% vs. cytology: 84%
Todenhöfer <i>et al.</i> (21)	2014	203	483	Patients undergoing surveillance of NMIBC	128	3, 7, 17, 9p21	Product labeling	UroVysion: 74% vs. cytology: 70%	UroVysion: 70% vs. cytology: 76%
Gomella <i>et al.</i> (22)	2017	284	415	Patients undergoing diagnosis or treatment of urothelial carcinoma	274	3, 7, 17, 9p21	Product labeling	UroVysion: 55% vs. cytology: 42%	UroVysion: 74% vs. cytology: 93%
Lavery <i>et al.</i> (23)	2017	129	129	Patients with bladder cancer before TURBT or RC	101	3, 7, 17, 9p21	Product labeling	UroVysion: 67% vs. cytology: 69%	UroVysion: 72% vs. cytology: 76%

BTA, bladder tumor antigen; NMIBC, non-muscle invasive bladder cancer; RC, radical cystectomy; TURBT, transurethral resection of bladder tumor; UC, urothelial carcinoma.

to detect upper tract UC in the follow-up of patients after cystectomy and concluded it was not suitable (39).

In summary, the sensitivity of UroVysion appears higher than that of urine cytology for diagnosing upper tract UC but indications for its routine use have not been established.

## Monitoring as follow-up

### *BCa surveillance without visible tumors*

Several reports exist of the usefulness of UroVysion in predicting BCa recurrence and progression. Yoder *et al.* (2007) reported a prospective study of patients under surveillance for BCa where recurrence was not shown. For 65% of a UroVysion-positive group, UC originated from bladder recurred within 29 months (40). Sarosdy *et al.* (2002) compared the predictive power of BCa recurrence between FISH, a BTA stat test, and cytology, and concluded overall sensitivities were 71% for FISH, 50% for the BTA stat test, and 26% for cytology (18). Seideman *et al.* (2015) also reported that in patients with a positive FISH result, BCa was more likely to recur regardless of a visible tumor (41). Kim *et al.* (2014) reported that being UroVysion-positive could predict the recurrence and progression of BCa in patients with negative cystoscopy and suspicious cytology results, and in those with a non-muscle invasive BCa negative result by cystoscopy (42). A prospective study comparing urinary biomarkers in a follow-up of patients with non-muscle invasive BCa with a low grade highlighted the superiority of UroVysion to cytology in terms of sensitivity and specificity (43). Consecutive UroVysion testing was reported as a more reliable method in predicting intravesical recurrence after transurethral resection of bladder tumor (TURBT) (15). In addition, these circumstances might change in the era of high-resolution endoscopes such as narrow band imaging or photodynamic diagnosis (15).

### *BCa surveillance after intravesical therapy*

UroVysion is also reported as a useful predictive test for anticipating BCa recurrence after intravesical therapy. Kipp *et al.* (2005) reported that monitoring using UroVysion was useful for predicting the progression of BCa to becoming muscle invasive in patients with intravesical therapy, including BCG and other intravesical therapies (44). Whitson *et al.* (2009) reported a UroVysion positive test after intravesical treatment was highly predictive of recurrence in a retrospective study (45). Regarding BCG intravesical treatment, Mengual *et al.* (2007) suggested that a UroVysion

result was useful for judging the need for adjuvant therapy in patients with a high-risk superficial BCa after an initial transurethral resection (46). Kamat *et al.* (2012) also reported a correlation between UroVysion results and the recurrence of BCa after BCG intravesical therapy; the timing of a positive test was important to predict recurrence (47). Freund *et al.* (2019) reported a significant correlation of recurrence in patients with intermediate- and high-risk UC originated from the bladder after intravesical BCG therapy and a positive UroVysion test 3 months following TURBT (35). A recent prospective, multicenter diagnostic trial followed previous results showing that a UroVysion positive result indicated a risk of recurrence (48). Though many studies suggest the effectiveness of a UroVysion test as a marker of BCa recurrence, some studies suggest the contrary. Savic *et al.* (2009) reported that a UroVysion test was superior to urine cytology only in the cases except for unequivocally positive cytology (49). They also concluded that a UroVysion test does not yield further information on clearly malignant cytology (49).

## Modification of UroVysion

Some studies tried to achieve better diagnostic accuracy than conventional UroVysion tests. Kipp *et al.* (2009) described the utility of counting abnormal cells, with the percentage of these a predictor of cancer recurrence and progression (50). Ferra *et al.* (2009) reported that less strict criteria dramatically improved the sensitivity of UroVysion instead of decreasing its specificity (29). The degree of chromosomal abnormalities, detected and assessed quantitatively by UroVysion, was reported as a predictor of the progression of non-muscle-invasive into muscle-invasive BCa (51). Ho *et al.* (2013) reported that specificity was improved by excluding chromosome 9 deletion; the use of UroVysion may allow the conservation of health resources and minimize trauma in patients by avoiding cystoscopy (52). Zhou *et al.* (2016) focused on tetrasomy and insisted this was a nonspecific finding that was frequently encountered and should be excluded from a polysomy classification (53). However, each study requires further investigation due to the small sample sizes used.

## Cost effectiveness of UroVysion

The costs associated with treating patients with BCa is highest among all cancers due to the monitoring and treatment required (54). The cost of diagnostic strategy in

BCa is expected to increase because of the arrival of new modalities. Physicians need to consider the cost effectiveness of each modality. The UroVysion test is a reasonable test for detecting BCa but shared positive test with urine cytology, which is a much cheaper test; its role is still questionable with regard to cost (29,31). Due to its high-cost limits and the fact that it is time consuming for use as a screening tool, a UroVysion test is not cost-effective (55). Kamat *et al.* (2011) examined the cost-effectiveness of using cystoscopy, in addition to urine cytology and urinary biomarkers, in their prospective study (56). They concluded that the cost per tumor detected using cystoscopy and UroVysion was much greater than that of cystoscopy on its own (\$19,111 vs. \$7,692) in following up of BCa; the addition of urinary biomarkers did not improve the detection of invasive disease. In another study, some experts were against using UroVysion in screening for hematuria due to its low cost-effectiveness and it being time-consuming (10,57,58).

### **The role of UroVysion in the era of diagnosis based on The Paris System on Reporting Urinary Cytology (TPS) since 2016**

For the long time, there was no standard diagnostic criteria of atypical urothelial cells universally available across different international institutions. For resolving this problem, TPS was proposed for pathological criteria using defined cytomorphologic classification for the diagnosis of urine cytology, and has been gradually prevailed. So far, several reports described that TPS could improve the reliability and accuracy of interpretations especially in urine cytology for high grade UC (HGUC). Based on the tendency, the number of articles presenting as to the efficacy of UroVysion using TPS. Miki *et al.* (2017) reviewed the UroVysion testing cases over a period of 6 years, and reclassified according to TPS, and reported that the sensitivity and specificity of UroVysion were 62.5% and 100% in the negative for HGUC groups (59). Recently, Tian *et al.* (2020) chronologically reviewed and retrospectively compared urine cytology diagnoses between pre-TPS and post-TPS, and described that all categories in urine cytology had superior correlation with UroVysion results by post-TPS, and positive UroVysion results increased from 87% to 93% in suspicious for HGUC category. By these results, it is hopeful that application of TPS could lead to significant reduction of atypical UC diagnosis, and potential cost effectiveness (60).

### **Future perspectives on UroVysion**

Recent advances in searching for genetic mutations have led to a paradigm shift in the treatment of cancers. In the case of observing genetic mutations in cells from urine samples, UroVysion has an important role to play in improving the management of cancer, especially UC. Several reports describe comparing or combining UroVysion with another type of genetic examination. Montalbo *et al.* (2020) reported that the sensitivities of UroVysion, the mutation analysis of fibroblast growth factor receptor 3 and telomerase reverse transcriptase, and urine-based gene expression analysis in the case of high-grade tumors were comparable but higher than that of cytology according to the Paris System for reporting urinary cytology reclassification (30). Chen *et al.* (2020) focused on urine DNA methylation to improve the sensitivity of detecting particularly low-grade and early-stage BCas compared with using cytology and UroVysion (61). Various urine-based examinations, including for genetic mutations, have been reported for decades but have not been found to be superior to UroVysion in detecting UC. Combining new types of examinations with UroVysion or using tailor-made examinations with various urine-based biomarkers are envisioned.

### **Limitations**

This narrative review has several limitations. First, quantitative analyses of the included studies were not conducted. Since the background of each study differs, differences in data between these studies need to be assessed statistically. Second, the inclusion/exclusion criteria used for the literature reviewed were ambiguous. To maintain the validity of the review, multiple authors were used to assess whether to include each article for this review.

### **Conclusions**

The accuracy of UroVysion tests can improve the diagnosis of UCs. Though various indications need to be considered by physicians when using UroVysion, cost concerns arise when this test is used blindly for screening the recurrence of UC. UroVysion has the potential to detect genetic mutations in cells from urine samples. Novel ways of improving or combining other types of pathological examinations with UroVysion are expected to provide new diagnostic tools in the treatment of UC.

## Acknowledgments

*Funding:* None.

## Footnote

*Provenance and Peer Review:* This article was commissioned by the Guest Editor (Kazutoshi Fujita) for the series “Urinary Biomarkers of Urothelial Malignancies” published in *Translational Andrology and Urology*. The article has undergone external peer review.

*Reporting Checklist:* The authors have completed the Narrative Review reporting checklist. Available at <http://dx.doi.org/10.21037/tau-20-1207>

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/tau-20-1207>). The series “Urinary Biomarkers of Urothelial Malignancies” was commissioned by the editorial office without any funding or sponsorship. The authors have no other conflicts of interest to declare.

*Ethical Statement:* the authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

- Richters A, Aben KKH, Kiemeny LALM. The global burden of urinary bladder cancer: an update. *World J Urol* 2020;38:1895-904.
- Soria F, Shariat SF, Lerner SP, et al. Epidemiology, diagnosis, preoperative evaluation and prognostic assessment of upper-tract urothelial carcinoma (UTUC). *World J Urol* 2017;35:379-87.
- Seklehner S, Remzi M, Fajkovic H, et al. Prospective Multi-institutional Study Analyzing Pain Perception of Flexible and Rigid Cystoscopy in Men. *Urology* 2015;85:737-41.
- Sathianathen NJ, Butaney M, Weight CJ, et al. Urinary Biomarkers in the Evaluation of Primary Hematuria: A Systematic Review and Meta-Analysis. *Bladder Cancer* 2018;4:353-63.
- Bhat A, Ritch CR. Urinary biomarkers in bladder cancer: Where do we stand? *Curr Opin Urol* 2019;29:203-9.
- Têtu B. Diagnosis of urothelial carcinoma from urine. *Mod Pathol* 2009;22:S53-9.
- Lokeshwar VB, Habuchi T, Grossman HB, et al. Bladder tumor markers beyond cytology: International Consensus Panel on bladder tumor markers. *Urology* 2005;66:35-63.
- Sokolova IA, Halling KC, Jenkins RB, et al. The development of a multitarget, multicolor fluorescence in situ hybridization assay for the detection of urothelial carcinoma in Urine. *J Mol Diagn* 2000;2:116-23.
- Bubendorf L. Multiprobe fluorescence in situ hybridization (UroVysion) for the detection of urothelial carcinoma - FISHing for the right catch. *Acta Cytol* 2011;55:113-9.
- Bubendorf L, Piaton E. UroVysion® multiprobe FISH in the triage of equivocal urinary cytology cases. *Ann Pathol* 2012;32:e52-6, 438-43.
- Miyake M, Owari T, Hori S, et al. Emerging biomarkers for the diagnosis and monitoring of urothelial carcinoma. *Res Rep Urol* 2018;10:251-61.
- Pesch B, Taeger D, Johnen G, et al. Screening for bladder cancer with urinary tumor markers in chemical workers with exposure to aromatic amines. *Int Arch Occup Environ Health* 2014;87:715-24.
- Zellweger T, Benz G, Cathomas G, et al. Multi-target fluorescence in situ hybridization in bladder washings for prediction of recurrent bladder cancer. *Int J Cancer* 2006;119:1660-5.
- Kipp BR, Sebo TJ, Griffin MD, et al. Analysis of Polyomavirus-Infected Renal Transplant Recipients' Urine Specimens. *Am J Clin Pathol* 2005;124:854-61.
- Ikeda A, Kojima T, Kawai K, et al. Risk for intravesical recurrence of bladder cancer stratified by the results on two consecutive UroVysion fluorescence in situ hybridization tests: a prospective follow-up study in Japan. *Int J Clin Oncol* 2020;25:1163-9.
- Sarosdy MF, Kahn PR, Ziffer MD, et al. Use of a Multitarget Fluorescence In Situ Hybridization Assay to Diagnose Bladder Cancer in Patients With Hematuria. *J Urol* 2006;176:44-7.
- Halling KC, King W, Sokolova IA, et al. A comparison

- of cytology and fluorescence in situ hybridization for the detection of urothelial carcinoma. *J Urol* 2000;164:1768-75.
18. Sarosdy MF, Schellhammer P, Bokinsky G, et al. Clinical evaluation of a multi-target fluorescent in situ hybridization assay for detection of bladder cancer. *J Urol* 2002;168:1950-4
  19. Laudadio J, Keane TE, Reeves HM, et al. Fluorescence in situ hybridization for detecting transitional cell carcinoma: implications for clinical practice. *BJU Int* 2005;96:1280-5.
  20. Caraway NP, Khanna A, Fernandez RL, et al. Fluorescence in situ hybridization for detecting urothelial carcinoma: a clinicopathologic study. *Cancer Cytopathol* 2010;118:259-68.
  21. Todenhöfer T, Hennenlotter J, Esser M, et al. Stepwise application of urine markers to detect tumor recurrence in patients undergoing surveillance for non-muscle-invasive bladder cancer. *Dis Markers* 2014;2014:973406.
  22. Gomella LG, Mann MJ, Cleary RC, et al. Fluorescence in situ hybridization (FISH) in the diagnosis of bladder and upper tract urothelial carcinoma: the largest single-institution experience to date. *Can J Urol* 2017;24:8620-6.
  23. Lavery HJ, Zaharieva B, McFaddin A, et al. A prospective comparison of UroVysion FISH and urine cytology in bladder cancer detection. *BMC Cancer* 2017;17:247.
  24. Hajdinjak T. UroVysion FISH test for detecting urothelial cancers: Meta-analysis of diagnostic accuracy and comparison with urinary cytology testing. *Urol Oncol* 2008;26:646-51.
  25. Moonen PMJ, Merckx GFM, Peelen P, et al. UroVysion Compared with Cytology and Quantitative Cytology in the Surveillance of Non-Muscle-Invasive Bladder Cancer. *Eur Urol* 2007;51:1275-80.
  26. Nagai T, Okamura T, Yanase T, et al. Examination of diagnostic accuracy of UroVysion fluorescence in situ hybridization for bladder cancer in a single community of Japanese hospital patients. *Asian Pac J Cancer Prev* 2019;20:1271-3.
  27. Daniely M, Rona R, Kaplan T, et al. Combined morphologic and fluorescence in situ hybridization analysis of voided urine samples for the detection and follow-up of bladder cancer in patients with benign urine cytology. *Cancer* 2007;111:517-24.
  28. Kojima T, Nishiyama H, Ozono S, et al. Clinical evaluation of two consecutive UroVysion fluorescence in situ hybridization tests to detect intravesical recurrence of bladder cancer: a prospective blinded comparative study in Japan. *Int J Clin Oncol* 2018;23:1140-7.
  29. Ferra S, Denley R, Herr H, et al. Reflex UroVysion testing in suspicious urine cytology cases. *Cancer* 2009;117:7-14.
  30. Montalbo R, Izquierdo L, Ingelmo-Torres M, et al. Urine cytology suspicious for urothelial carcinoma: Prospective follow-up of cases using cytology and urine biomarker-based ancillary techniques. *Cancer Cytopathol* 2020;128:460-9.
  31. Bonberg N, Taeger D, Gawrych K, et al. Chromosomal instability and bladder cancer: the UroVysion(TM) test in the UroScreen study. *BJU Int* 2013;112:E372-E382.
  32. Yu J, Xiong H, Wei C, et al. Utility of fluorescence in situ hybridization analysis for detecting upper urinary tract-urothelial carcinoma. *J Cancer Res Ther* 2017;13:647.
  33. Marín-Aguilera M, Mengual L, Ribal MJ, et al. Utility of Fluorescence In Situ Hybridization as a Non-invasive Technique in the Diagnosis of Upper Urinary Tract Urothelial Carcinoma. *Eur Urol* 2007;51:409-15.
  34. Mian C, Mazzoleni G, Vikoler S, et al. Fluorescence in situ hybridisation in the diagnosis of upper urinary tract tumours. *Eur Urol* 2010;58:288-92.
  35. Freund JE, Liem EIML, Savci-Heijink CD, et al. Fluorescence in situ hybridization in 1 mL of selective urine for the detection of upper tract urothelial carcinoma: a feasibility study. *Med Oncol* 2018;36:10.
  36. Su X, Hao H, Li X, et al. Fluorescence in situ hybridization status of voided urine predicts invasive and high-grade upper tract urothelial carcinoma. *Oncotarget* 2017;8:26106-11.
  37. Todenhöfer T, Hennenlotter J, Esser M, et al. Combined application of cytology and molecular urine markers to improve the detection of urothelial carcinoma. *Cancer Cytopathol* 2013;121:252-60.
  38. Sassa N, Iwata H, Kato M, et al. Diagnostic Utility of UroVysion Combined With Conventional Urinary Cytology for Urothelial Carcinoma of the Upper Urinary Tract. *Am J Clin Pathol* 2019;151:469-78.
  39. Fernández MI, Parikh S, Grossman HB, et al. The role of FISH and cytology in upper urinary tract surveillance after radical cystectomy for bladder cancer. *Urol Oncol* 2012;30:821-4.
  40. Yoder BJ, Skacel M, Hedgepeth R, et al. Reflex UroVysion testing of bladder cancer surveillance patients with equivocal or negative urine cytology: A prospective study with focus on the natural history of anticipatory positive findings. *Am J Clin Pathol* 2007;127:295-301.
  41. Seideman C, Canter D, Kim P, et al. Multicenter evaluation of the role of UroVysion FISH assay in surveillance of patients with bladder cancer: does FISH positivity anticipate recurrence? *World J Urol* 2015;33:1309-13.

42. Kim PH, Sukhu R, Cordon BH, et al. Reflex fluorescence in situ hybridization assay for suspicious urinary cytology in patients with bladder cancer with negative surveillance cystoscopy. *BJU Int* 2014;114:354-9.
43. Benderska-Söder N, Hovanec J, Pesch B, et al. Toward noninvasive follow-up of low-risk bladder cancer - Rationale and concept of the UroFollow trial. *Urol Oncol* 2020;38:886-95.
44. Kipp BR, Karnes RJ, Brankley SM, et al. Monitoring intravesical therapy for superficial bladder cancer using fluorescence in situ hybridization. *J Urol* 2005;173:401-4.
45. Whitson J, Berry A, Carroll P, et al. A multicolour fluorescence in situ hybridization test predicts recurrence in patients with high-risk superficial bladder tumours undergoing intravesical therapy. *BJU Int* 2009;104:336-9.
46. Mengual L, Marín-Aguilera M, José Ribal M, et al. Clinical Utility of Fluorescent in situ Hybridization for the Surveillance of Bladder Cancer Patients Treated with Bacillus Calmette-Guérin Therapy. *Eur Urol* 2007;52:752-9.
47. Kamat AM, Dickstein RJ, Messetti F, et al. Use of fluorescence in situ hybridization to predict response to bacillus Calmette-Guérin therapy for bladder cancer: Results of a prospective trial. *J Urol* 2012;187:862-7.
48. Lotan Y, Inman BA, Davis LG, et al. Evaluation of the Fluorescence In Situ Hybridization Test to Predict Recurrence and/or Progression of Disease after bacillus Calmette-Guérin for Primary High Grade Nonmuscle Invasive Bladder Cancer: Results from a Prospective Multicenter Trial. *J Urol* 2019;202:920-6.
49. Savic S, Zlobec I, Thalmann GN, et al. The prognostic value of cytology and fluorescence in situ hybridization in the follow-up of nonmuscle-invasive bladder cancer after intravesical Bacillus Calmette-Guérin therapy. *Int J Cancer* 2009;124:2899-904.
50. Kipp BR, Tanasescu M, Else TA, et al. Quantitative fluorescence in situ hybridization and its ability to predict bladder cancer recurrence and progression to muscle-invasive bladder cancer. *J Mol Diagn* 2009;11:148-54.
51. Kocsmár I, Pajor G, Gyöngyösi B, et al. Development and Initial Testing of a Modified UroVysion-Based Fluorescence In Situ Hybridization Score for Prediction of Progression in Bladder Cancer. *Am J Clin Pathol* 2020;153:274-84.
52. Ho CCK, Tan WP, Pathmanathan R, et al. Fluorescence-in-situ-hybridization in the surveillance of urothelial cancers: Can use of cystoscopy or ureteroscopy be deferred? *Asian Pac J Cancer Prev* 2013;14:4057-9.
53. Zhou AG, Liu Y, St. Cyr M, et al. Role of tetrasomy for the diagnosis of urothelial carcinoma using UroVysion fluorescent in situ hybridization. *Arch Pathol Lab Med* 2016;140:552-9.
54. Botteman MF, Pashos CL, Redaelli A, et al. The Health Economics of Bladder Cancer: A Comprehensive Review of the Published Literature. *Pharmacoeconomics* 2003;21:1315-30.
55. Zhou AG, Owens CL, Cosar EF, et al. Clinical implications of current developments in genitourinary pathology. *Arch Pathol Lab Med* 2013;137:887-93.
56. Kamat AM, Karam JA, Grossman HB, et al. Prospective trial to identify optimal bladder cancer surveillance protocol: reducing costs while maximizing sensitivity. *BJU Int* 2011;108:1119-23.
57. Nieder AM, Soloway MS, Herr HW. Should We Abandon the FISH Test? *European Urology* 2007;51:1469-71.
58. Budman LI, Kassouf W, Steinberg JR. Biomarkers for detection and surveillance of bladder cancer. *Can Urol Assoc J* 2008;2:212-21.
59. Miki Y, Neat M, Chandra A. Application of The Paris System to atypical urine cytology samples: correlation with histology and UroVysion® FISH. *Cytopathology* 2017;28:88-95.
60. Tian W, Shore KT, Shah RB. Significant reduction of indeterminate (atypical) diagnosis after implementation of The Paris System for reporting urinary cytology: A single-institution study of more than 27,000 cases. *Cancer Cytopathol* 2021;129:114-20.
61. Chen X, Zhang J, Ruan W, et al. Urine DNA methylation assay enables early detection and recurrence monitoring for bladder cancer. *J Clin Invest* 2020;130:6278-89.

**Cite this article as:** Nagai T, Naiki T, Etani T, Iida K, Noda Y, Shimizu N, Isobe T, Nozaki S, Okamura T, Ando R, Kawai N, Yasui T. UroVysion fluorescence in situ hybridization in urothelial carcinoma: a narrative review and future perspectives. *Transl Androl Urol* 2021;10(4):1908-1917. doi: 10.21037/tau-20-1207